

Symplectic Gravity Models in Four, Three and Two Dimensions

Oleg Kechkin and Maria Yurova

Nuclear Physics Institute,

Moscow State University,

Moscow 119899, RUSSIA,

e-mail: kechkin@cdfs.npi.msu.su

(October 1996)

Abstract

A class of the $D = 4$ gravity models describing a coupled system of n Abelian vector fields and the symmetric $n \times n$ matrix generalizations of the dilaton and Kalb–Ramond fields is considered. It is shown that the Pecci–Quinn axion matrix can be entered and the resulting equations of motion possess the $Sp(2n, R)$ symmetry in four dimensions. The stationary case is studied. It is established that the theory allows a σ –model representation with a target space which is invariant under the $Sp[2(n+1), R]$ group of isometry transformations. The chiral matrix of the coset $Sp[2(n+1), R]/U(n+1)$ is constructed. A Kähler formalism based on the use of the Ernst $(n+1) \times (n+1)$ complex symmetric matrix is developed. The stationary axisymmetric case is considered. The Belinsky–Zakharov chiral matrix depending on the original field variables is obtained. The Kramer–Neugebauer transformation, which algebraically maps the original variables into the target space ones, is presented.

PACS numbers: 04.20.Jb, 04.50.+h

Typeset using REVTeX

I. INTRODUCTION

Last years much attention had been attracted by the study of symmetries for gravity models arising as the low energy limit of superstring theory [1]– [7]. This activity was motivated by the problems of superstring theory, as well as by the remarkable results obtained for the Einstein and Einstein–Maxwell theories in three and two dimensions. Namely, it had been established the chiral matrix formulations for the latter theories in the stationary case, and their complete integrability in the stationary axisymmetric case [8]– [21].

Recently it was shown that the simplest string gravity model, the Einstein–Maxwell–Dilaton–Axion (EMDA) system, can be represented as the 2×2 matrix generalization of the pure Einstein theory [22]– [23]. It means that one can exchange the complex symmetric 2×2 matrix, describing the EMDA theory in three dimensions, by the Ernst potential of the Einstein theory to obtain this last one from the former system [22]. Thus, the mentioned matrix (the Ernst matrix potential) generalizes the Ernst potential to the EMDA case, and both the Einstein and EMDA theories belong to the same class of gravity models. In [23] it was shown that the established analogy is even the more complete one, and can be naturally prolonged to the stationary axisymmetric case. It had been shown, that the EMDA system, as the Einstein theory, allows both the target space representation and the formulation based on the use of the original field components. These two formulations are differentially related in three dimensions and can be algebraically mapped one into another in two dimensions [23]. This map is known as the Kramer–Neugebauer transformation for the stationary axisymmetric Einstein equations [24]; its existence is an important property of the theory [25].

The formal properties of the EMDA theory do not actually depend on the Ernst matrix potential dimension and it is possible to consider a series of similar gravity models in three and two dimensions. All these models exist as the chiral ones and possess the remarkable features of the Einstein theory. But first of all one must give the answer to the question what kind of the four–dimensional theories being reduced to three and two dimensions lead

to such models.

In this paper we present such a class of four-dimensional theories. Any theory generalizes the EMDA system by the following way: here the dilaton and Kalb–Ramond fields become the $n \times n$ symmetric matrices, and instead of the Maxwell field one has the column of n Maxwell fields (thus, the EMDA theory corresponds to the case of $n = 1$).

It is shown that one can introduce the Pecci–Quinn axion matrix for an arbitrary $n \geq 1$ and that the resulting equations of motion possess the $Sp(2n, R)$ hidden symmetry. This fact generalizes the one for the EMDA theory which $Sp(2, R) \sim SL(2, R)$ symmetry is known as the S –duality [26]–[27]. Also it is shown that the model admits the discrete symmetry transformation which had been named as the “strong–weak coupling duality” for the low energy heterotic string theory with moduli fields in four dimensions [3].

In the next section we consider the stationary case and reduce the four-dimensional model with an arbitrary n to three dimensions. We use an approach which was firstly formulated by Israel and Wilson for the Einstein–Maxwell theory in [28], and recently had been generalized to the EMDA theory in [29]. It is shown that the system under consideration allows the σ –model representation with a symmetric target space, which is a natural matrix generalization of the EMDA one.

Then, the Kähler formalism based on the use of the $(n + 1) \times (n + 1)$ Ernst matrix potential is developed. It is shown that the motion equations become the matrix-valued Ernst equation. The chiral $2(n + 1) \times 2(n + 1)$ matrix constructed on the real and imaginary parts of the Ernst matrix potential is presented. This matrix belongs to the coset $Sp[2(n + 1), R]/U(n + 1)$, and the established chiral formulation possesses the $Sp[2(n + 1), R]$ group of symmetry transformations. The corresponding transformations for the Ernst matrix potential have the “matrix-valued $SL(2, R)$ form” and directly generalize the transformations for the Einstein theory. Also it is shown that the $Sp(2n, R)$ symmetry, corresponding to the one of the four-dimensional motion equations, exists as the subgroup of the complete $Sp[2(n + 1), R]$ symmetry group. At the end of the section the discrete transformation, which is the analogy of the strong–weak coupling duality for the low energy heterotic string

theory in three dimensions [4], is presented.

In the following section we demonstrate some new features of the model, which arise in the stationary axisymmetric case. It is established that the theory allows the alternative Ernst-like formulation using two real $(n+1) \times (n+1)$ matrices, which are the nondualized (i.e., depended on the original non-target space field components) analogies of the Ernst matrix and its complex conjugation. These two real matrices are used for the construction of the new $2(n+1) \times 2(n+1)$ chiral matrix which coincides with the Belinsky–Zakharov one for the case of the Einstein theory (see also [23] for the EMDA theory case). This new matrix undergoes the same $Sp[2(n+1), R]$ transformations as the coset chiral matrix under the action of the isometry group. The above mentioned real Ernst-like matrix potentials transform as their complex Ernst analogies.

At the end of the paper we present the Kramer–Neugebauer map for an arbitrary n . It turned out that this transformation has an especially simple form in terms of the real and complex pairs of the Ernst matrix potentials.

II. FOUR-DIMENSIONAL EQUATIONS

In this paper we consider a class of gravity models with the action

$$^4S = \int d^4x |g|^{\frac{1}{2}} \{-R + Tr[\frac{1}{2}(\partial p p^{-1})^2 - p F F^T + \frac{1}{3}(p H)^2]\}, \quad (1)$$

where R is the Ricci scalar for the metric $g_{\mu\nu}$, ($\mu = 0, \dots, 3$); $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ and

$$H_{\mu\nu\lambda} = \partial_\mu B_{\nu\lambda} - \frac{1}{2}(A_\mu F_{\nu\lambda}^T + F_{\nu\lambda} A_\mu^T) + \text{cyclic}. \quad (2)$$

Here the symmetric $n \times n$ matrix p is constructed on the scalar field components ($p = e^{-2\phi}$ for the EMDA case, where ϕ is the dilaton field), $B_{\mu\nu}$ is the symmetric $n \times n$ matrix which contains the antisymmetric tensor Kalb–Ramond fields (i.e., $B_{\mu\nu}^T = B_{\mu\nu}$ and $B_{\mu\nu} = -B_{\nu\mu}$), and A_μ is the $n \times 1$ column of the Abelian vector fields. This action describes the Einstein and EMDA theories in the cases of $n = 0$ and $n = 1$ respectively and provides their natural

generalization for an arbitrary n . The models of such type arise in the low energy limit of the heterotic string theory after the compactification of extra dimensions on a torus [1]– [7].

As it has been done for the EMDA theory and for the string gravity with moduli fields, one can establish an alternative formulation of the problem which is connected with the dualization of the axion field $H_{\mu\nu\lambda}$ in four dimensions. Namely, using the first equation of the Euler–Lagrange ones,

$$\nabla^\mu(pH^{\mu\nu\lambda}p) = 0, \quad (3)$$

$$\nabla_\mu(pF^{\mu\nu}) + \frac{1}{2}pH^{\mu\nu\lambda}F_{\mu\nu} = 0, \quad (4)$$

$$\nabla J^p + pFF^T - \frac{2}{3}(pH)^2 = 0, \quad (5)$$

$$R_{\mu\nu} = Tr[\frac{1}{2}J_\mu^p J_\nu^p - 2p(F_{\mu\lambda}F^{T\nu\lambda} - \frac{1}{4}g_{\mu\nu}FF^T) + pH_{\mu\lambda\sigma}pH_\nu^{\lambda\sigma} - \frac{1}{3}g_{\mu\nu}(pH)^2] \quad (6)$$

where $J^p = (\nabla p)p^{-1}$, it is possible to introduce the symmetric pseudoscalar matrix q ,

$$H^{\mu\nu\lambda} = \frac{1}{2}E^{\mu\nu\lambda\sigma}p^{-1}q_{,\sigma}p^{-1} \quad (7)$$

generalizing the Pecci–Quinn axion field. It satisfies the relation

$$\nabla J^q - J^p J^q + p\tilde{F}F^T = 0, \quad (8)$$

where $J^q = (\nabla q)p^{-1}$ and $\tilde{F}^{\mu\nu} = \frac{1}{2}E^{\mu\nu\lambda\sigma}F_{\lambda\sigma}$, which together with Eqs. (4)–(6), rewritten using Eq. (7) as

$$\nabla_\mu(pF^{\mu\nu} + q\tilde{F}^{\mu\nu}) = 0, \quad (9)$$

$$\nabla J^p + (J^q)^2 + pFF^T = 0, \quad (10)$$

$$R_{\mu\nu} = Tr[\frac{1}{2}(J_\mu^p J_\nu^p + J_\mu^q J_\nu^q) - 2p(F_{\mu\lambda}F^{T\nu\lambda} - \frac{1}{4}g_{\mu\nu}FF^T)] \quad (11)$$

form the complete system for the action

$${}^4S = \int d^4x |g|^{\frac{1}{2}} \{-R + Tr[\frac{1}{2}((J^p)^2 + (J^q)^2) - pFF^T - q\tilde{F}F^T]\}. \quad (12)$$

The action of this type firstly was considered by Breitenlohner, Gibbons and Maison in [31], where some remarkable properties of the model had been established.

The main purpose of this paper is to show a complete formal analogy between the series of models describing by the action (1), and the Einstein theory in three and two dimensions. However, the simplest representative of the series, which possesses all the typical properties in four dimensions, is the EMDA theory. Namely, let us establish the symmetries of the theory for an arbitrary $n \geq 1$. To do it one can introduce the complex variables

$$z = q + ip, \quad (13)$$

$$\mathcal{F} = \frac{1}{2}(F + i\tilde{F}). \quad (14)$$

Then the set of motion equations obtains the form:

$$\nabla_\mu(z\mathcal{F}^{\mu\nu} - \bar{z}\bar{\mathcal{F}}^{\mu\nu}) = 0, \quad (15)$$

$$\nabla J^z - J^z(J^z - \bar{J}^z) - \frac{i}{2}(z - \bar{z})\bar{\mathcal{F}}\mathcal{F}^+ = 0, \quad (16)$$

$$R_{\mu\nu} = Tr\{2J_{(\mu}^z\bar{J}_{\nu)}^z + i(z - \bar{z})(\mathcal{F}_{\mu\lambda}\mathcal{F}^+_{\nu}{}^\lambda + \bar{\mathcal{F}}_{\mu\lambda}\mathcal{F}^T_{\nu}{}^\lambda)\}, \quad (17)$$

where $J^z = \nabla z(z - \bar{z})^{-1}$ (see also [26] for the EMDA case).

It is not difficult to prove using a straightforward calculation that these equations are invariant under the transformation

$$z \rightarrow s^T(z^{-1} + l)^{-1}s + r, \quad \mathcal{F} \rightarrow s^{-1}(I + lz)\mathcal{F}, \quad (18)$$

where the constant matrices $r^T = r$, $l^T = l$ and s are real. The former formula from (18) also can be rewritten using the following $2n \times 2n$ matrices

$$m = \begin{pmatrix} p^{-1} & p^{-1}q \\ qp^{-1} & p + qp^{-1}q \end{pmatrix} \quad (19)$$

(see the similar matrix for the Einstein–Dilaton–Axion theory in [32])

$$g = \begin{pmatrix} (s^T)^{-1} & (s^T)^{-1}r \\ l(s^T)^{-1} & s + l(s^T)^{-1}r \end{pmatrix}, \quad (20)$$

as

$$m \rightarrow g^T m g. \quad (21)$$

The above introduced matrix g satisfies the relation $g^T j g = j$, where

$$j = \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix}, \quad (22)$$

and I is the unit matrix, so that $g \in Sp(2n, R)$ (also from the relations $m^T j m = j$ and $m^T = m$ it follows that $m \in Sp(2n, R)/U(n)$, see the third chapter for details). Thus, the motion equations possess the $Sp(2n, R)$ symmetry for an arbitrary $n \geq 1$.

Not all the symplectic transformations (21) can be represented in the Gauss decomposition form (20). For example, one can prove that $j \in Sp(2n, R)$, but only the limit procedure defined by the matrices $l = -r = s \rightarrow \infty$ allows to obtain the matrix j from Eq. (20). The corresponding map is $m \rightarrow m^{-1}$; and using complex variables one has:

$$z \rightarrow -z^{-1}, \quad \mathcal{F} \rightarrow z\mathcal{F}. \quad (23)$$

It is not difficult to prove that this transformation is actually the symmetry transformation for the equations (15)–(17). The map (23) generalizes the strong–weak coupling duality of the EMDA theory to the case of an arbitrary n . It has the same form as the transformation established for the low energy heterotic string theory with moduli in four dimensions [3].

It is easy to see that only the $GL(n, R)$ subgroup of the complete $Sp[2(n+1), R]$ group, defined by a matrix s , preserves the action (12) which can be rewritten as

$$^4S = \int d^4x |g|^{\frac{1}{2}} \{-R + Tr[2J^z \bar{J}^z + i(z\mathcal{F}\mathcal{F}^T - \bar{z}\bar{\mathcal{F}}\mathcal{F}^+)]\}. \quad (24)$$

The situation is the same one as for the EMDA theory which possesses the $Sp(2, R) \sim SL(2, R)$ symmetry only for the motion equations [26]–[27]. This symmetry is connected

with the notable S -duality for the low energy heterotic string theory. The subgroup $SL(2, Z)$ of the group $SL(2, R)$ is an exact symmetry for the heterotic string in four dimensions [3].

Also one can see that the discussed models do not possess any analogy with the T -duality transformations [3], thus the established $Sp(2n, R)$ symmetry is the single non-trivial one.

III. REDUCTION TO THREE DIMENSIONS

In this section we study the stationary case using the classical approach of Israel and Wilson [28]. This method also had been explored for the EMDA theory analysis [29]; its application to the string gravity with moduli fields one can find in [4].

First of all, it is convenient to parametrize the four-dimensional line element as

$$ds_4^2 = g_{\mu\nu} dx^\mu dx^\nu = f(dt - \omega_m dx^m)^2 - f^{-1} h_{mn} dx^m dx^n, \quad (25)$$

where $m = 1, 2, 3$. (In this section all vector variables, as well as the operator ∇ , will be related with the three-metric h_{mn}).

Then, the $\mu = m$ component of Eq. (9) allows to introduce the magnetic potential u according to

$$pF^{mn} + q\tilde{F}^{mn} = \frac{1}{\sqrt{2}} f E^{mnk} u_{,k}, \quad (26)$$

while for the electric potential v we put

$$F_{m0} = \frac{1}{\sqrt{2}} v_{,m}. \quad (27)$$

Using the relations [28] $F^{0m} = \omega_n F^{mn} + h^{mn} F_{n0}$ and $F_{mn} = f^{-2} h_{mk} h_{nl} F^{kl} + 2F_{0[m} \omega_{n]}$, it is possible to rewrite the $\mu = 0$ component of Eq. (9) as

$$\nabla[f^{-1}(p\nabla v - qp^{-1}(\nabla u - q\nabla v))] = -f^{-2} \vec{\tau} \nabla u, \quad (28)$$

where $\vec{\tau} = -f^2 \nabla \times \vec{\omega}$. The second electromagnetic equation can be obtained from the Bianchi identity $F_{[mn,k]} = 0$; the result is:

$$\nabla[f^{-1}p^{-1}(\nabla u - q\nabla v)] = f^{-2}\vec{\tau}\nabla v. \quad (29)$$

The equations (8) and (10) transform to

$$\nabla J^q - J^p J^q - f^{-1}[(\nabla u - q\nabla v)\nabla v^T + p\nabla v(\nabla u - q\nabla v)^T p^{-1}] = 0, \quad (30)$$

$$\nabla J^p + (J^p)^2 + f^{-1}[p\nabla v\nabla v^T - (\nabla u - q\nabla v)(\nabla u - q\nabla v)^T p^{-1}] = 0. \quad (31)$$

Then, the $({}^m_0)$ -component of the Einstein equations (11) means the existence of the rotational potential χ

$$\nabla\chi + v^T\nabla u - u^T\nabla v = \vec{\tau}, \quad (32)$$

which accordingly with the definition of $\vec{\tau}$ satisfies

$$\nabla[f^{-2}(\nabla\chi + v^T\nabla u - u^T\nabla v)] = 0; \quad (33)$$

while the $({}_{00})$ -component of Eq. (11) leads to the following equation for the function f :

$$\begin{aligned} & f\nabla^2 f - (\nabla f)^2 + (\nabla\chi + v^T\nabla u - u^T\nabla v)^2 - \\ & f[\nabla v^T p\nabla v + (\nabla u - q\nabla v)^T p^{-1}(\nabla u - q\nabla v)] = 0. \end{aligned} \quad (34)$$

Finally, the $({}^{mn})$ -component of Eq. (11) after the lowering of the three-dimensional indexes takes the form:

$$\begin{aligned} {}^3R_{mn} = & \frac{1}{2}Tr[J_m^p J_n^p + J_m^q J_n^q] + \frac{1}{2}f^{-2}[f_{,m}f_{,n} + \vec{\tau}_m\vec{\tau}_n] \\ & - f^{-1}[v_{,m}^T p v_{,n} + (u_{,m} - qv_{,m})^T p^{-1}(u_{,n} - qv_{,n})], \end{aligned} \quad (35)$$

where ${}^3R$ is constructed using the metric h_{mn} and $\vec{\tau}_m$ is given by Eq. (32).

One can prove that Eqs. (28)–(31) and (33)–(35) are the Euler–Lagrange ones for the action

$$\begin{aligned} {}^3S = & \int d^3x h^{\frac{1}{2}} \{ - {}^3R + \frac{1}{2}Tr[(J^p)^2 + (J^q)^2] + \frac{1}{2}f^{-2}[(\nabla f)^2 + (\nabla\chi + v^T\nabla u - u^T\nabla v)^2] \\ & - f^{-1}[\nabla v^T p\nabla v + (\nabla u - q\nabla v)^T p^{-1}(\nabla u - q\nabla v)] \}. \end{aligned} \quad (36)$$

Hence, this system allows the sigma-model representation in the stationary case.

This result contains the one for the EMDA theory [29]–[30], which admits some compact matrix formulations. To establish them for an arbitrary n , it is convenient to introduce the complex potentials

$$\Phi = u - zv, \quad (37)$$

$$\mathcal{E} = if - \chi + v^T \Phi,$$

which together with the matrix z , entered before (13), form the set of Ernst-like potentials [33]–[34] for this problem (see also [22] for the EMDA case). These potentials can be combined into the following $(n+1) \times (n+1)$ symmetric matrix

$$E = \begin{pmatrix} \mathcal{E} & \Phi^T \\ \Phi & -z \end{pmatrix}, \quad (38)$$

which provides the Kähler representation of the model. Namely, it is easy to check that

$$^3S = \int d^3x h^{\frac{1}{2}} \{-^3R + 2Tr(J^E \bar{J}^E)\}, \quad (39)$$

where $J^E = \nabla E (E - \bar{E})^{-1}$ and the 2-form $\mathcal{K} = 2iTr(\mathcal{J}^E \wedge \bar{\mathcal{J}}^E)$, corresponding to the target space metric $dl^2 = 2Tr(\mathcal{J}^E \bar{\mathcal{J}}^E)$ with $\mathcal{J}^E = dE (E - \bar{E})^{-1}$, is exact. The Kähler formulations of the Einstein and Einstein–Maxwell theories are well known in the literature [16]. The another examples of the Kähler systems one can find in [35].

The equations following from the action (39)

$$\nabla J^E = J^E (J^E - \bar{J}^E), \quad (40)$$

$$^3R_{mn} = 2Tr(J_{(m}^E \bar{J}_{n)}^E) \quad (41)$$

coincide with the Ernst ones [33] in the case of $n = 0$. Thus, it is natural to name the matrix function E as the “matrix Ernst potential” for an arbitrary n . This Ernst-like representation for the model provides the above mentioned complete formal analogy between it and the Einstein theory in the stationary case.

Then, using the real and imaginary parts of $E = Q + iP$,

$$P = \begin{pmatrix} f - v^T p v & -v^T p \\ -p v & -p \end{pmatrix}, \quad Q = \begin{pmatrix} v^T w - \chi & w^T \\ w & -q \end{pmatrix}, \quad (42)$$

where $w = u - qv$, one can rewrite the action (39) as

$$^3S = \int d^3x h^{\frac{1}{2}} \{ -^3R + \frac{1}{2} Tr[(J^P)^2 + (J^Q)^2] \}, \quad (43)$$

with $J^P = (\nabla P)P^{-1}$ and $J^Q = (\nabla Q)P^{-1}$.

Now let us establish an alternative matrix formulation of the problem based on the use of the original non-target space variables. One can see that the first equation of motion corresponding to the action (43),

$$\nabla J^Q - J^P J^Q = 0, \quad (44)$$

$$\nabla J^P + (J^Q)^2 = 0, \quad (45)$$

$$^3R_{mn} = \frac{1}{2} Tr[J_m^P J_n^P + J_m^Q J_n^Q], \quad (46)$$

being written as $\nabla[P^{-1}(\nabla Q)P^{-1}] = 0$, ensures the compatibility condition for the relation

$$\nabla \times \vec{\Omega} = P^{-1}(\nabla Q)P^{-1} \quad (47)$$

which defines the vector matrix $\vec{\Omega}$. Then, using the relations (7), (26), (32) and the definition of $\vec{\tau}$ one obtains the explicit form of this matrix:

$$\vec{\Omega} = \begin{pmatrix} \vec{\omega} & -\sqrt{2}(\vec{A} + A_0 \vec{\omega})^T \\ -\sqrt{2}(\vec{A} + A_0 \vec{\omega}) & 2\vec{B} + (\vec{A} + A_0 \vec{\omega})A_0^T + A_0(\vec{A} + A_0 \vec{\omega})^T \end{pmatrix}, \quad (48)$$

where $B_m = 2B_{0m}$. It is easy to see that the matrix $\vec{\Omega}$ consists of the original field variables from Eqs. (1) and (2) which had been named as “nondualized” ones in [23] for the case of the EMDA theory.

The remaining components of the Kalb–Ramond field $B_{mn} \equiv \frac{1}{2} E_{mnk} C^k$ satisfy the non-dynamical equation

$$\begin{aligned}\nabla \vec{C} = & \{\nabla[\vec{B} \times \vec{\omega}] + [\vec{B} - (\vec{A} + A_0\vec{\omega})A_0^T - A_0(\vec{A} + A_0\vec{\omega})^T]\nabla \times \vec{\omega} \\ & + (\vec{A} + A_0\vec{\omega})\nabla \times (\vec{A} + A_0\vec{\omega})^T\} + \nabla \times (\vec{A} + A_0\vec{\omega})(\vec{A} + A_0\vec{\omega})^T,\end{aligned}\quad (49)$$

as it follows from Eqs. (7) and the t -independence of the q .

Then, it is easy to see from Eqs. (47) that the matrices P and $\vec{\Omega}$ satisfy the relation $\nabla \times [P\nabla \times (\vec{\Omega})P] = 0$ which also can be rewritten using the matrix current $J^{\vec{\Omega}} = P\nabla \times \vec{\Omega}$ as

$$\nabla \times J^{\vec{\Omega}} - J^{\vec{\Omega}} \times J^P = 0. \quad (50)$$

This relation together with Eqs. (45) and (46), expressed in terms of the matrices P and $\vec{\Omega}$,

$$\nabla J^P + (J^{\vec{\Omega}})^2 = 0, \quad (51)$$

$${}^3R_{mn} = \frac{1}{2}Tr[J_m^P J_n^P + J_m^{\vec{\Omega}} J_n^{\vec{\Omega}}], \quad (52)$$

form the set of motion equations for the action

$${}^3S = \int d^3x h^{\frac{1}{2}}(-{}^3R + \frac{1}{2}Tr[(J^P)^2 - (J^{\vec{\Omega}})^2]); \quad (53)$$

thus these matrices provide an alternative Lagrange formulation of the theory under consideration.

To establish the symmetry group of the system, it is convenient to introduce the $2(n+1) \times 2(n+1)$ symmetric matrix

$$M = \begin{pmatrix} P^{-1} & P^{-1}Q \\ QP^{-1} & P + QP^{-1}Q \end{pmatrix}, \quad (54)$$

which possesses the symplectic property

$$M^T J M = J, \quad \text{where} \quad J = \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix}. \quad (55)$$

Then the action (43) obtains the form

$${}^3S = \int d^3x h^{\frac{1}{2}} \{-{}^3R + \frac{1}{4}Tr[(J^M)^2]\}, \quad (56)$$

where $J^M = \nabla M M^{-1}$, and the corresponding equations of motion are:

$$\nabla J^M = 0, \quad (57)$$

$${}^3R_{mn} = \frac{1}{4}Tr[J_m^M J_n^M]. \quad (58)$$

It is known that in the case of $n = 0$ (for the Einstein theory) a matrix M belongs to the coset $Sp(2, R)/U(1)$ (or, equivalently, to the $SL(2, R)/O(2)$) [12]. Also, it has been established that for the EMDA theory one has $M \in Sp(4, R)/U(2)$ [22], [36]. Hence, it is natural to suppose that $M \in Sp[2(n+1), R]/U(n+1)$ for an arbitrary n .

This supposition really takes place and the local isomorphism can be established as follows. One can see that a matrix M remains symplectic and symmetric under the transformation

$$M \rightarrow G^T M G \quad (59)$$

with an arbitrary symplectic matrix G . It means that the group of the isometry transformations for the action (56) is isomorphic to $Sp[2(n+1), R]$. The set of the corresponding generators $\Gamma = G - I$ can be obtained from (59); the result is:

$$\Gamma = \begin{pmatrix} -\Gamma_S^T & \Gamma_R \\ \Gamma_L & \Gamma_S \end{pmatrix}, \quad (60)$$

where $(n+1) \times (n+1)$ infinitesimal matrices Γ_R and Γ_L are symmetric, and Γ_S is an arbitrary matrix of the same dimension. Then, to obtain the coset, one must remove all the antisymmetric generators from eq. (60), i. e., to exclude the set of generators which is defined by the matrix

$$\tilde{\Gamma} = \begin{pmatrix} \tilde{\Gamma}_S & \tilde{\Gamma}_R \\ -\tilde{\Gamma}_R & \tilde{\Gamma}_S \end{pmatrix}, \quad (61)$$

where $\tilde{\Gamma}_S^T = -\tilde{\Gamma}_S$ and $\tilde{\Gamma}_R^T = \tilde{\Gamma}_R$.

From the other hand, the generators u of the group $U(n+1)$ can be found from the relation $U^+U = I$ in the infinitesimal case of $U = I + u$. The result is:

$$u = u_1 + iu_2, \quad (62)$$

where $u_1^T = -u_1$ and $u_2^T = u_2$ are the real $(n+1) \times (n+1)$ matrices. Now one can prove using a straightforward calculation of the commutators for the corresponding generators that the waiting isomorphism can be defined as

$$\tilde{\Gamma}_S \equiv u_1, \quad \tilde{\Gamma}_R \equiv u_2. \quad (63)$$

Now let us establish the action of the $Sp[2(n+1), R]$ group of the isometry transformations on the Ernst matrix potential E . One can see that the matrix

$$G = \begin{pmatrix} (S^T)^{-1} & (S^T)^{-1}R \\ L(S^T)^{-1} & S + L(S^T)^{-1}R \end{pmatrix}, \quad (64)$$

with $R^T = R$ and $L^T = L$ possesses the symplectic property and describes all the belonging to $Sp[2(n+1), R]$ transformations which can be continuously connected with the identical one. It is easy to show that the corresponding transformation for the matrix E is

$$E \rightarrow S^T(E^{-1} + L)^{-1}S + R. \quad (65)$$

Thus, the matrix E undergoes the “matrix-valued $SL(2, R)$ ” transformations which are the matrix generalizations of the $SL(2, R)$ ones for the Einstein theory. Namely, the transformation defined by the matrix L is the matrix generalization of the Ehlers one [37], while the matrices S and R define the transformations which generalize the rescaling and gauge shift correspondingly for the stationary Einstein system [12].

The transformation (65) has the same form as the transformation for the matrix z in Eq. (18). It is not difficult to prove that one can identify the $Sp(2n, R)$ subgroup of the complete group $Sp[2(n+1), R]$, defined by the matrices

$$L = \begin{pmatrix} 0 & 0 \\ 0 & l \end{pmatrix}, \quad S = \begin{pmatrix} 1 & 0 \\ 0 & s \end{pmatrix}, \quad R = \begin{pmatrix} 0 & 0 \\ 0 & r \end{pmatrix}, \quad (66)$$

where $l^T = l$ and $r^T = r$, with the $Sp(2n, R)$ symmetry transformations (18) of the motion equations (15)–(17) in the stationary case.

As an example of a symplectic transformation which can not be represented in the form of (64), one can take the matrix J . It is easy to see that the corresponding map is $M \rightarrow M^{-1}$ and its complex representation has the form $E \rightarrow -E^{-1}$. This transformation can be obtained from the decomposition formula (64) using the limit procedure as it has been done in the four-dimensional case. It also provides an analogy with the strong–weak coupling duality transformation of the low energy heterotic string theory with moduli in three dimensions [4].

At the end of this section we want to note that all the transformations discussed here preserve both the actions and the corresponding sets of motion equations.

IV. TWO-DIMENSIONAL CASE

In this section we consider some additional properties of the theory which arise in the stationary and axisymmetric case.

Here one can parametrize the three-dimensional line element in the Lewis–Papapetrou form:

$$ds_3^2 = e^{2\gamma}(d\rho^2 + dz^2) + \rho^2 d\varphi^2. \quad (67)$$

Then Eq. (57) transforms to

$$\nabla(\rho J^M) = 0 \quad (68)$$

which is the Euler–Lagrange equation for the action

$$^2S = \frac{1}{4} \int d\rho dz \rho \text{Tr}[(J^M)^2]; \quad (69)$$

while the Einstein equations (58) become the relations defining the function γ :

$$\begin{aligned}\gamma_{,z} &= \frac{\rho}{4} \text{Tr}[J_\rho^M J_z^M], \\ \gamma_{,\rho} &= \frac{\rho}{8} \text{Tr}[(J_\rho^M)^2 - (J_z^M)^2]\end{aligned}\tag{70}$$

(the operator ∇ is connected with the flat metric δ_{ab} in this section).

The remarkable fact is that one can obtain the another chiral matrix for the stationary axisymmetric equations of the problem. To make it, we firstly consider the formalism based on the use of the matrices P and Q . It is easy to see that the φ -independent equations of motion (44)–(45) take the form

$$\nabla(\rho J^Q) - \rho J^P J^Q = 0,\tag{71}$$

$$\nabla(\rho J^P) + \rho (J^Q)^2 = 0,\tag{72}$$

and can be derived from the action

$$^2S = \frac{1}{2} \int d\rho dz \rho \text{Tr}[(J^P)^2 + (J^Q)^2].\tag{73}$$

Also, from Eq. (46) one obtains:

$$\begin{aligned}\gamma_{,z} &= \frac{\rho}{2} \text{Tr}[J_\rho^P J_z^P + J_\rho^Q J_z^Q], \\ \gamma_{,\rho} &= \frac{\rho}{4} \text{Tr}[(J_\rho^P)^2 - (J_z^P)^2 + (J_\rho^Q)^2 - (J_z^Q)^2].\end{aligned}\tag{74}$$

The equation (71), being written as $\nabla[\rho P^{-1}(\nabla Q)P^{-1}] = 0$, provides the compatibility condition for the relation

$$\nabla\Omega = \rho P^{-1}(\tilde{\nabla}Q)P^{-1}\tag{75}$$

which defines the symmetric matrix Ω (here, accordingly with [12], $\tilde{\nabla}_\rho = \nabla_z$ and $\tilde{\nabla}_z = -\nabla_\rho$).

Then, using a straightforward calculation one obtains that

$$\Omega = \vec{\Omega}_\varphi.\tag{76}$$

It is easy to see that the compatibility condition for the relation (75), rewritten as $\nabla Q = -\rho^{-1}P(\tilde{\nabla}\Omega)P$, and the equation (72), expressed in terms of the matrices P and Ω ,

$$\nabla(\rho^{-1}J^\Omega) + \rho^{-1}J^\Omega J^P = 0, \quad (77)$$

$$\nabla(\rho^{-1}J^P) + \rho^{-1}(J^\Omega)^2 = 0, \quad (78)$$

form the Euler–Lagrange set of equations for the action

$$^2S = \frac{1}{2} \int d\rho dz Tr[\rho(J^P)^2 - \rho^{-1}(J^\Omega)^2], \quad (79)$$

where $J^\Omega = P\nabla\Omega$. Also one can see from Eq. (49), that $\nabla\vec{C} = 0$ and, fixing a gauge, it is possible to put $\vec{C} = 0$ without lose of generality. Then, for the function γ one has:

$$\begin{aligned} \gamma_{,z} &= \frac{1}{2} Tr[\rho J_\rho^P J_z^P - \rho^{-1} J_\rho^\Omega J_z^\Omega], \\ \gamma_{,\rho} &= \frac{1}{4} Tr[\rho((J_\rho^P)^2 - (J_z^P)^2) + \rho^{-1}((J_\rho^\Omega)^2 - (J_z^\Omega)^2)]. \end{aligned} \quad (80)$$

The new chiral matrix N can be defined as follows

$$N = \begin{pmatrix} P & -P\Omega \\ -\Omega P & \Omega P\Omega - \rho^2 P^{-1} \end{pmatrix}. \quad (81)$$

It satisfies the equation

$$\nabla[\rho J^N] = 0, \quad (82)$$

where $J^N = \nabla N N^{-1}$, while the relations for the function γ are:

$$\begin{aligned} \Gamma_{,z} &= \frac{\rho}{4} Tr[J_\rho^N J_z^N], \\ \Gamma_{,\rho} &= \frac{\rho}{8} Tr[(J_\rho^N)^2 - (J_z^N)^2], \end{aligned} \quad (83)$$

with

$$\Gamma = \gamma - \frac{1}{2} \ln | \det P | + \frac{n+1}{2} \ln \rho. \quad (84)$$

It is easy to check that the introduced symmetric matrix N satisfies the non-group relation

$$N J N = -\rho^2 J \quad (85)$$

which is equivalent to the normalization condition $\det N = -\rho^2$ in the case of the Einstein theory (see also [23] for the EMDA case).

An analogy between the model under consideration and the Einstein theory allows to establish one additional property of the system. Namely, the relation (75) provides the differential correspondence between the formulations of the problem based on the use of the dualized matrix Q and the original one Ω . The remarkable fact is that also there is a pure algebraical map

$$P \rightarrow \rho P^{-1}, \quad Q \rightarrow i\Omega, \quad (86)$$

which directly transforms the equations (71)–(72) into the ones (77)–(78). The corresponding transformation for the function γ is:

$$e^{2\gamma^\Omega} = \frac{\rho^{\frac{n+1}{2}}}{|\det P|} e^{2\gamma^Q}, \quad (87)$$

where γ^Ω is connected with matrices P and Ω , and γ^Q is related with the formulation using P and Q . The map (86) coincides with the Kramer–Neugebauer transformation in the cases of $n = 0$ [24] and $n = 1$ [23] and can naturally preserve this name for an arbitrary n .

Now let us establish the action of the $Sp[2(n+1), R]$ group on the matrices P and Ω . To make it one can introduce the matrices

$$E_\pm = \Omega \pm \rho P^{-1}. \quad (88)$$

It is easy to check that the matrix N remains symmetric and preserves the property (85) under the transformation $N \rightarrow G^T N G$ with an arbitrary symplectic matrix G . Then, in the case of the matrix G , possessing the decomposition property (64), one has:

$$E_\pm \rightarrow S^T (E_\pm^{-1} + L)^{-1} S + R. \quad (89)$$

Thus, the real nondualized matrix variables $E_{\pm}$ transform under the $Sp[2(n+1), R]$ group of transformations as the matrix Ernst potentials E and $\bar{E}$.

The entered matrices $E_{\pm}$, being the analogy for the Ernst ones E and $\bar{E}$, provide also the compact Lagrange formulation of the problem. Namely, one can check that the action (79) can be rewritten as

$$^2S = -2 \int d\rho dz \rho Tr[J^{E+} J^{E-}], \quad (90)$$

where $J^{E\pm} = (E_{\pm} - E_{\mp})^{-1} \nabla E_{\pm} - \frac{1}{2} \nabla \rho$.

Also one can see that the above established Kramer–Neugebauer map has an especially simple form in terms of the Ernst matrices $E, \bar{E}$ and their real analogies E_+, E_- :

$$E \rightarrow iE_+, \quad \bar{E} \rightarrow iE_-. \quad (91)$$

V. CONCLUSION

In this paper we have considered a class of gravity models with the Einstein theory as the first representative, the EMDA theory as the second one, etc. It is shown that all the models have the same formal properties which are the natural matrix generalizations of the ones for the Einstein theory in three and two dimensions. Namely, the substitution $P \rightarrow f$, $Q \rightarrow \chi$ and $\vec{\Omega} \rightarrow \vec{\omega}$ (or $\Omega \rightarrow \omega$ in the stationary axisymmetric case) directly transforms all the relations for $n \geq 1$ to the Einstein theory relations. An analogy established here is even a more complete one, and it is possible to develop the Hauser–Ernst formalism [17], [19] and to construct the Geroch group [8]–[9] in the stationary axisymmetric case for the model with an arbitrary n .

ACKNOWLEDGMENTS

We would like to thank our colleagues from the Nuclear Physics Institute for an encouraging relation to our work.

REFERENCES

- [1] J. Maharana and J.H. Schwarz, Nucl. Phys. **B390** (1993) 3.
- [2] J.H. Schwarz and A. Sen, Nucl. Phys. **B411** (1993) 35.
- [3] A. Sen, Int. J. Mod. Phys. **A9** (1993) 3707.
- [4] A. Sen, Nucl. Phys. **B434** (1995) 179.
- [5] A. Sen, Nucl. Phys. **B447** (1995) 62.
- [6] J. Maharana, Phys. Rev. Lett. **75** (1995) 208.
- [7] A.K. Biswas, A. Kumar and K. Ray, Nucl. Phys. **B453** (1995) 181.
- [8] R. Geroch, J. Math. Phys. **12** (1971) 918.
- [9] R. Geroch, J. Math. Phys. **13** (1972) 394.
- [10] V.A. Belinsky and V.E. Zakharov, Zh. Exp. Teor. Fiz. **75** (1978) 1955 [Sov. Phys. JETP **48** (1978) 985].
- [11] V.A. Belinsky and V.E. Zakharov, Zh. Exp. Teor. Fiz. **77** (1979) 3 [Sov. Phys. JETP **50** (1979) 1].
- [12] W. Kinnersley, J. Math. Phys. **18** (1977) 529.
- [13] W. Kinnersley and D. M. Chitre, J. Math. Phys. **18** (1977) 1538.
- [14] W. Kinnersley and D. M. Chitre, J. Math. Phys. **19** (1978) 1926.
- [15] W. Kinnersley and D. M. Chitre, J. Math. Phys. **19** (1978) 2037.
- [16] P.O. Mazur, Acta Phys. Pol. **14** (1983) 219.
- [17] I. Hauser and F.J. Ernst, Phys. Rev. **D20** (1979) 362.
- [18] I. Hauser and F.J. Ernst, Phys. Rev. **D20** (1979) 1783.

- [19] I. Hauser and F.J. Ernst, J. Math. Phys. **21** (1980) 1126.
- [20] I. Hauser and F.J. Ernst, J. Math. Phys. **21** (1980) 1418.
- [21] C.M. Cosgrove, J. Math. Phys. **22** (1981) 2624.
- [22] D.V. Gal'tsov and O.V. Kechkin, Phys. Lett. **B361** (1995) 52.
- [23] O. Kechkin and M. Yurova, “Kramer-Neugebauer transformation for Einstein-Maxwell-dilaton-axion theory”, to be published in Phys. Rev. **D54** (1996); report hep-th/9604175.
- [24] G. Neugebauer and D. Kramer, Ann. Phys. (NY) **24** (1969) 62.
- [25] P. Breitenlohner and D. Maison, Ann. Poincare **46** (1987) 215.
- [26] A. Shapere, S. Trivedi and F. Wilczec, Mod. Phys. Lett. **A6** (1991) 2677.
- [27] R. Kallosh and T. Ortin, Phys. Rev. **D48** (1993) 742.
- [28] W. Israel and G. A. Wilson, J. Math. Phys. **13** (1972) 865.
- [29] D.V. Gal'tsov, A.A. Garcia and O.V. Kechkin, J. Math. Phys. (1995) 5023.
- [30] D.V. Gal'tsov and O.V. Kechkin, Phys. Rev. **D50** (1994) 7394.
- [31] P. Breitenlohner, D. Maison and G. Gibbons, Commun. Math. Phys. **120** (1988) 295.
- [32] I. Bakas, Nucl. Phys. **B428** (1994) 374.
- [33] F.J. Ernst, Phys. Rev. **167** (1968) 1175.
- [34] F.J. Ernst, Phys. Rev. **168** (1968) 1415.
- [35] P. Fre, Nucl. Phys. Proc. Suppl. **45BC** (1996) 59.
- [36] D.V. Gal'tsov and O.V. Kechkin, Phys. Rev. **D54** (1996) 1656.
- [37] J. Ehlers, in “Les Theories de la Gravitation” (CNRS, Paris, 1959).